CUMULATIVE INDEX 2001

Volume 32

January	FOOT AND ANKLE TRAUMA, pages 1–218
April	ADVANCED TECHNIQUES IN THE MANAGEMENT OF WRIST TRAUMA, pages 219–379
July	REPAIR OF ATHLETIC SHOULDER INJURIES, pages 381–542
October	INSTABILITY AFTER MAJOR JOINT REPLACEMENT, pages 543–722

Note: Page numbers of article and issue titles are in **boldface** type.

Abductor digiti minimi muscle, as flap for lateral calcaneal defects, 155–157
Abductor hallucis muscle, as flap for calcaneal fractures with medial ankle or heel wound, 157–158
Abductor tension, increasing, for treatment of hip instability, 606
Acceleration vs. deceleration injury, in throwers with SLAP lesions, 436–437
Acetabular component repositioning, for treatment of hip instability, 599–600
Acetabular implants design, change of, for treatment of hip instability, 600–603
Acromioclavicular joint, in arthroscopy for full-thickness rotator cuff tears, 504
  sprained, in athletes, 394
Acromioplasty, for full-thickness rotator cuff tears, 504
Adductor tenotomy, for treatment of hip instability, 608
Allograft bone, as bone graft substitute for unstable distal radius fractures, 342
Allograft bone matrix, as bone graft substitute for unstable distal radius fractures, 342–343
Allograft-implant composites, for distal femur reconstruction, 706–707
  for pelvic reconstruction, 700
  for proximal femur reconstruction, 703
  for proximal tibia reconstruction, 707
Amputation, after complex foot and ankle trauma. See *Foot and ankle trauma, complex, salvage or amputation after.*
Ankle, anatomy and biomechanics, 79–80
  vascular anatomy, 136–141
Ankle fractures, complications, 88–89
    in diabetic patients, **113–133**
      authors' clinical series, 129–130
      Charcot arthropathy, 119–123
      clinical questions, about displaced ankle fractures, 123–127
        about nondisplaced ankle fractures, 123
        about reducing complications from surgery for displaced ankle fractures, 127–129
        about role for fracture healing adjuncts, 129
        about type of fixation for osteopenic bone, 129
        clinical series, 113–119
          delayed fracture healing, 115–116
          wound healing, 116–119
        summary, 130–131
    pathology, 83
    physical examination, 83
    principles of fixation, lateral malleolus, 87–88
      medial malleolus, 88
      posterior malleolus, 88
    radiographic evaluation, 80–83
    specific, management of, bimalleolar fractures and equivalents, 85
      isolated lateral malleolus fracture, 84
      isolated medial malleolus fracture, 84–85
      Maisonneuve fractures and syndesmosis injuries, 85–86
      nonsurgical vs. surgical, 84
      open fractures, 86–87
      trimalleolar fractures, 85
    summary, 89
    and syndesmosis injuries, **79–90**
Anterior superior instability with rotator cuff tearing: SLAC lesion, **457–461**
  clinical examination, 457–458
  diagnostic studies, 458
  discussion, 459–461
  materials and methods, 457
  results, 459
  surgical findings, 458–459
Arthrodesis, instability in total knee arthroplasty and, 623–624
  for pelvic reconstruction, 700–702
Arthroplasty. See *specific types of arthroplasty.*
Arthroscopic Bankart repair with Suretac device for traumatic anterior shoulder instability in athletes, **411–421**
  anatomy of shoulder stability, capsule, 412
    glenoid labrum, 411–412
  complications, 418
  discussion, 418–419

Arthroscopic (*Continued*)
patient selection, arthroscopic evaluation, 413
examination under anesthesia, 412–413
general considerations, 412
radiographic evaluation, 412
postoperative care, 415–416
results, 416–417
surgical technique, arthroscopic evaluation, 414
drilling glenoid and tensioning soft tissue, 414–415
glenoid and labrum preparation, 414
implant device and instrumentation, 413–414
placement of implant, 415
thermal capsulorrhaphy, 416
Arthroscopic repair of partial-thickness rotator cuff tears and SLAP lesions in professional baseball players, **443–456**
instability, 444
internal impingement, 443
MRI imaging, 445–446
results, 454–455
rotator cuff tears, 445
SLAP lesions, 444–445
subacromial impingement, 445
surgical technique, arthroscopic examination, 447
debridement, 448
electrothermal capsule repair, 450, 452
examination under anesthesia, 446
knot tying principles, 450
order of repair, 447
portal placement, 446–447
position and traction, 446
postoperative protocol, 453–454
rotator cuff repair, 452–453
superior labrum repair, 448–450
Arthroscopic rotator cuff repairs, all, **501–510**
discussion, 509–510
full-thickness rotator cuff tears, acromioclavicular joint, 504
acromioplasty, 504
anchor placement, 505
anchor selection, 505
cuff mobilization, 503–504
instruments, 508
knot tying, 507
repair site preparation, 504–505
suture placement and passing, 505, 507
suture selection, 505
tear classification, 503
visualization, 503
indications, 501
operative technique, inspection of glenohumeral joint, 502–503
portals, 502
positioning, 501–502
partial-thickness rotator cuff tears, 503
postoperative management, 508
results, 508–509
Arthroscopic sliding knot, "new," **523–524**
Arthroscopy, for dynamic scaphoid instability, 236–239
monopolar electrothermal, for treatment of shoulder instability in athlete, **511–521**
applications of thermal stabilization of shoulder, 514
clinical results, 518–519
complications and failures, 519–520
methods of thermal energy delivery, 512–514
postoperative care, multidirectional instability, 518
posterior instability, 517–518
unidirectional anterior instability, 517
scientific background, 511–512
surgical technique, 514–517
for scaphoid fractures, 251
history, 255–256
for SLAP lesions, 437–438
treating distal radius fractures with, **307–315**
indications and timing, 309
methods of articular evaluation, 307–309
outcomes, 313–314
postoperative care, 313
summary, 314
technique, 309–313
portals, 310–311
tips and tricks for fracture reduction, 311–313
treatment of associated injuries, 313
wrist distraction, 310
for triangular fibrocartilage complex injuries, **279–294**
anatomy, 279–281
biomechanics, 281
classification, 284–286
diagnosis, diagnostic modalities, 283–284
differential diagnosis, 282
history, 282
physical examination, 282
for irreparable lesions, 291–292
postoperative care, 292
for reparable radial-sided lesions, 289–291
for reparable ulnar-sided lesions, 286, 288–289
results, 292
summary, 292–293
treatment, nonoperative, 286
open surgical repair, 286, 287
ulnar recession, 286
Athletic shoulder injuries, repair of, all arthroscopic rotator cuff repairs, **501–510**
anatomy, pathophysiology, and biomechanics of shoulder instability, **381–391**
anterior superior instability with rotator cuff tearing: SLAC lesion, **457–461**
arthroscopic Bankart repair with Suretac device for traumatic anterior shoulder instability in athletes, **411–421**
arthroscopic repair of partial-thickness rotator cuff tears and SLAP lesions in professional baseball players, **443–456**
coracoid impingement syndrome, rotator interval reconstruction, and biceps tenodesis in, **485–493**
differential diagnosis of shoulder injuries in sports, **393–398**
evaluation and management of shoulder instability in elite overhead thrower, **423–430**
issue on, **381–542**
monopolar electrothermal arthroscopy for treatment of shoulder instability in athlete, **511–521**
"new" arthroscopic sliding knot, **523–524**
posterior instability, **463–473**
shoulder rehabilitation strategies, guidelines, and practice, **525–536**
SLAP lesions in overhead athlete, **431–441**
technique for repair of subscapularis tendon, **495–500**
traumatic anterior shoulder instability, **399–409**
treatment of athlete with multidirectional shoulder instability, **475–484**

Bankart repair, arthroscopic. See *Arthroscopic Bankart repair with Suretac device for traumatic anterior shoulder instability in athletes.*
suture anchor placement and, for traumatic anterior shoulder instability, 404–406
Biceps lesions, in overhead athlete, 488–489
aftercare, 492
anatomy and pathophysiology, 489
surgical technique, 491–492
treatment, 489–491
Bimalleolar fractures, and equivalents, 85
Bioactive glass, as bone graft substitute for unstable distal radius fractures, 346–349
Bohler's angle, and wound healing after open reduction and internal fixation of calcaneal fractures, 187–192
Bone graft, and alternatives, role of, in unstable distal radius fracture treatment, **337–351**
approved bone graft substitutes, 339–340
basic science of remodeling, 338, 342
bone graft substitutes, allograft bone, 342
allograft bone matrix, 342–343
bioactive glass, 346–347
bovine bone–derived ceramic, 347–348
calcium phosphate cement, 345–346
calcium sulfate, 347
ceramics and ceramic composites, 343–344
collagen mineral composite graft, 344
coralline hydroxyapatite, 344–345
bone graft substitutes in development, 341
bioactive glass, 348–349
calcium phosphate, 348
calcium phosphate cements, 348
osteoinductive materials, 349
measurement of osteoinduction and osteoconduction, 342
problems with existing materials, 348
summary, 349–350
vascularized. See *Vascularized bone grafts.*
Bone-retinaculum-bone reconstruction, of scapholunate ligament injuries, **241–246**
discussion, 244–245
indications, 241–242
outcomes, 244
postoperative care, 243–244
surgical technique, 242–243
Bovine bone–derived ceramic, as bone graft substitute for unstable distal radius fractures, 347–348

Calcaneal fractures, **35–51**
authors' preferred method of treatment, 49
classification, 39, 41
complications, 49–50
conclusions, 50
definitive management, extra-articular fractures, 44–45
intra-articular fractures, 45–47
history, 35
initial management, 41, 43
initial presentation, 41
mechanism of injury, 37–39
in multiply injured patient, 198–199
or osteomyelitis, soft tissue reconstruction for, **135–170**
approach to acute calcaneus fracture, 146
arterial blood supply to lateral calcaneal area, development of angiosome concept, 135–136
lateral calcaneal exposure, using angiosome concept, 141, 146
vascular anatomy of lateral ankle and heel, 136–141
management of lateral calcaneal wounds, approach to acute lateral calcaneal wound, 150–151
approach to acutely infected lateral calcaneal wound, 151
approach to chronic lateral calcaneal wound, 152
postdebridement wound management, 152
role of debridement, 146, 150
reconstruction of acute or chronic lateral calcaneal wound, delayed primary closure, 153
local flaps, 153, 155
microsurgical free flaps, 158, 165
pedicled flaps, 155–158
skin graft, 153
scope of the problem, 135
summary, 165–168
postoperative management, 49
radiographic examination, 43–44
salvage procedures, 50
surgical approach, lateral, 47–48
medial, 48–49
use of ring fixators in, 210–212
wound-healing risk factors after open reduction and internal fixation of, **187–192**
discussion, 191–192
materials and methods, 187–189
operative technique, 189
results, 189–191
Calcaneus, anatomy, 35–37
Calcific tendinopathy, in athletes, 397
Calcium phosphate, as bone graft substitute for unstable distal radius fractures, 348
cements, as bone graft substitute for unstable distal radius fractures, 345–346, 348
Calcium sulfate, as bone graft substitute for unstable distal radius fractures, 347
Capsule, electrothermal repair of, in professional baseball players, 450, 452
pathoanatomy of, 412
Carpal kinematics, normal and abnormal, advances in in vivo measurement of, **219–231**
abnormal carpal motion: SLIL injured versus uninjured kinematics, 228–229
carpal bone motion patterns, effect of scapholunate interosseous ligament tears, 220
row versus column, 219–220
need for measuring in vivo carpal kinematics, 221
normal carpal motion, capitate kinematics, 224–226
row versus column, 222–224
scaphoid and lunate kinematics, 226–228
overview of methods for markerless bone registration, 221–222
purpose, 219
summary and major findings, 229
summary of previous measurement methodologies, 220–221
Carpus, acute fracture–dislocation of, complications, 304
diagnosis, 296–297
and management of, **295–305**
epidemiology, 295–296
mechanism of injury, 296
results, 301–302
summary, 304
treatment, 298–301
anatomy, 295
Ceramics and ceramic composites, as bone graft substitute for unstable distal radius fractures, 343–344

Charcot joints, instability in total knee arthroplasty and, 624
Charcot neuropathy, in diabetic patients with ankle fractures, 119–123
  ring fixators for, 209
Clavicle fractures, in athletes, 394
Closed reduction, and cast, for treatment of instability after total elbow arthroplasty, 689
  as definitive treatment of hip instability, 597–598
Collagen mineral composite graft, as bone graft substitute for unstable distal radius fractures, 344
Component revision, for treatment of instability after total elbow arthroplasty, 692–693
Coracoid impingement, anatomy and pathophysiology, 485
  in overhead athlete, aftercare, 486
    clinical presentation, physical examination, and diagnosis, 486
    treatment, 486
Coracoid impingement syndrome, rotator interval reconstruction, and biceps tenodesis in overhead athlete, **485–493**
Coralline hydroxyapatite, as bone graft substitute for unstable distal radius fractures, 344–345
Cuboid bone, dislocations of, 24
  fractures of, 22–24
  stress fractures of, 24–25
Cuneiform bones, fractures and dislocations of, 29–30

Debridement, role of, in lateral calcaneal wounds, 146, 150
Diabetic patients, complications of ankle fractures in, **113–133**
  authors' clinical series, 129–130
  Charcot arthropathy, 119–123
  clinical questions, about displaced ankle fractures, 123–127
    about nondisplaced ankle fractures, 123
    about reducing complications from surgery for displaced ankle fractures, 127–129
    about role for fracture healing adjuncts, 129
    about type of fixation for osteopenic bone, 129
  clinical series, delayed fracture healing, 115–116
    wound healing, 116–119
  summary, 130–131
Dislocation, of cuboid bone, 24
  of cuneiform bone, 29–30
  of first metatarsophalangeal joint, 7–8
  of hip, preoperative planning to prevent, **579–586**
    decreasing incidence of dislocation, component design and size, 582–583
    component positioning, 583–584
    operative factors, 581
    patient risk factors, 579–581
    postoperative treatment, 584
  interphalangeal, 8–9
  vs. subluxation, in hip instability, 593–594
Distal femur, instability after major tumor resection in, 705–707
Distal radial ulnar joint deterioration, treatment of, 365–367
Distal radial ulnar joint replacement, prostheses used for,
  Herbert technique, 369–370
  Scheker technique, 370–373
  Swanson technique, 367–369
  uHead prosthesis, 370
Distal radioulnar joint, posttraumatic unstable, reconstruction of, **353–363**
  discussion, 361–362
  history and physical examination, 354
  indications and contraindications, 354–355
  operative technique, 355–360
  radiographic evaluation, 354
  results, 360–361
Distal radius fractures, five-pin external fixation and early range of motion for, **329–335**
  discussion, 331–334
  summary, 334–335
  technique, 330–331
  limited incision open techniques for management of, **317–327**
  augmentation of fracture stability, 318–319
  biomechanical analysis, 319
  case studies, 322–326
  contraindications, 322, 324
  evolution of treatment, 318
  predictors of outcome, 317–318
  principles and methods of fragment-specific fixation, 319–322
    exposure, 320
    implant application, 320–321
  summary, 324–325
  treating, with arthroscopic assistance, **307–315**
    indications and timing, 309
    methods of articular evaluation, 307–309
    outcomes, 313–314
    postoperative care, 313
    summary, 314
    technique, 309–313
      portals, 310–311
      tips and tricks for fracture reduction, 311–313
      treatment of associated injuries, 313
      wrist distraction, 310
  unstable, 337–338
    role of bone graft and alternatives in treatment of, **337–351**
    approved bone graft substitutes, 339–340
      allograft bone, 342
      allograft bone matrix, 342–343
      bioactive glass, 346–347
      bovine bone–derived ceramic, 347–348
      calcium phosphate cement, 345–346
      calcium sulfate, 347
      ceramics and ceramic composites, 343–344
      collagen mineral composite graft, 344
      coralline hydroxyapatite, 344–345
    basic science of remodeling, 338, 342
    bone graft substitutes in development, 341
      bioactive glass, 348–349
      calcium phosphate, 348
      calcium phosphate cements, 348
      osteoinductive materials, 349
    measurement of osteoinduction and osteoconduction, 342
    problems with existing materials, 348
    summary, 349–350
Distal ulnar prosthetic replacement, **365–376**
  discussion, 374–375
  Herbert technique, 369–370
  Scheker technique, 370–373
  summary, 375
  Swanson technique, 367–369
  uHead prosthesis, 370
Dorsal distal radius and ulna, vascular anatomy of, 264–267

Electrothermal arthroscopy, monopolar. See *Monopolar electrothermal arthroscopy for treatment of shoulder instability in athlete.*
Electrothermal capsule repair, in professional baseball players, 450, 452
Endoprostheses, for distal femur reconstruction, 705
for pelvic reconstruction, 699
for proximal femur reconstruction, 703
for proximal tibia reconstruction, 707
Essential joints, as consideration for salvage vs. amputation after complex foot and ankle trauma, 183
Extra-articular deformity, correcting, for preventing instability in total knee arthroplasty, 623

Fascia lata plasty, for treatment of hip instability, 607–608
Femoral and tibial component rotation, intraoperatively determining, during primary or revision total knee arthroplasty, 640–641
preoperatively determining, in malfunctioning total knee arthroplasty, 641–644
Femoral bone resection, for preventing instability in total knee arthroplasty, 616–619
Femoral component, increasing offset or height of, for treatment of hip instability, 606
reorientation of, for treatment of hip instability, 603
Femur, distal, instability after major tumor resection in, 705–707
proximal, instability after major tumor resection in, 703–705
total replacement of, for tumor, instability after, 708–709
Fifth metatarsal. See *Metatarsal(s), fifth.*
Five-pin external fixation and early range of motion, for distal radius fractures, **329–335**
discussion, 331–334
summary, 334–335
technique, 330–331
Fixed flexion deformity, treating, for preventing instability in total knee arthroplasty, 622–623
Flaps, local, for acute or chronic lateral calcaneal wound, 153, 155
microsurgical free, for soft tissue reconstruction for calcaneal fractures or osteomyelitis, 158, 165
pedicled, for acute or chronic lateral calcaneal wound, 155–158
Flexion and extension gap, balancing, for preventing instability in total knee arthroplasty, 620–621
Flexion instability, of knee, 632–634
Foot and ankle trauma, ankle fractures and syndesmosis injuries, **79–90**
complex, applications of ring fixators in, **205–214**
conclusions, 212
historical perspective, 205
reasons for using, 205–206
specific fractures, calcaneal fractures, 210–212
foot fractures, 212
neuroarthropathic (Charcot) fractures, 209
talus fractures, 209–210
tibial plafond fractures, 206–209
salvage or amputation after, **181–186**
considerations influencing, distal injuries, 183
essential joints, 183
fracture, 182–183
free vascularized tissue transfers, 183–185
level of amputation, 185–186
muscle balancing, 186
nerve transection, 182
soft tissue injury, 181–182
structural integrity, 182
vascular injury, 182
zone of injury, 182
summary, 186
complications of ankle fractures in diabetic patients, **113–133**
foot compartment syndrome, **103–111**
fractures of base of fifth metatarsal, **171–180**
fractures of calcaneus, **35–51**
fractures of tarsal bones, **21–33**
injuries to toes and metatarsals, **1–10**
issue on, **1–218**
outcome of foot injuries in multiply injured patients, **193–204**
recent developments and advances in diagnosis and treatment of injuries to tarsometatarsal joint, **11–20**
soft tissue reconstruction for calcaneal fractures or osteomyelitis, **135–170**
talar body fractures, **65–77**
talar neck fractures, **53–64**
treatment of pilon fractures, **91–102**
wound-healing risk factors after open reduction and internal fixation of calcaneal fractures, **187–192**
Foot compartment syndrome, **103–111**
considerations of time and energy, 106
diagnosing, in children, 107–108
clinical symptoms, 107
differential diagnosis, 107
open fracture and foot compartment syndrome, 107
experimentally induced, 103
foot compartment anatomy, 104–105
from foot injuries in multiply injured patient, 196
incidence, 105–106
nontraumatic causes, 106
pathophysiology, **103–111**
summary, 110
treatment, authors' preferred algorithm for, 109
authors' preferred method of operative management of complications, 110
natural history of foot compartment syndrome, 108
nonoperative management of complications, 109
surgical, 108–109
vs. nonsurgical, 109
Foot injuries, outcome of, in multiply injured patients, **193–204**
authors' review, 199–202
compartment syndrome, 196
general treatment principles, 195–197
high-energy injuries, 194–195
initial management of trauma patient, 193–194
investigations, 197–198
missed or delayed diagnosis, 194
reasons for, 202
specific injuries, calcaneus, 198–199
Lisfranc, 199
metatarsal, 199
midfoot, 199
talus, 198
timing of surgery, 197
vascular compromise, 196
Forefoot fractures, use of ring fixators in, 212
Fracture–dislocation, acute, of carpus, anatomy, 295
complications, 304
diagnosis, 296–297
and management, **295–305**
epidemiology, 295–296
mechanism of injury, 296

Fracture–dislocation (*Continued*)
results, 301–302
summary, 304
treatment, 298–301
Fractures. See *specific fractures*.
Free vascularized tissue transfers, as consideration for salvage vs. amputation after complex foot and ankle trauma, 183–185

Glenohumeral arthroplasty, factors determining mobility after, 662–663
factors determining stability after, 663–664
unstable, clinical syndromes, 664–666
evaluation, 666
management and prevention, 666–668
Glenohumeral dislocations, in athletes, 394–395
Glenohumeral joint, posterior instability of, **463–473**
authors' preferred operative technique, 467–470
discussion, 471–472
evaluation, 464–465
pathophysiology, 463–464
rehabilitation, 470
results, 470–471
treatment, 465–467
Glenoid labrum, pathoanatomy of, 411–412

Heel, vascular anatomy of, 136–141
Hemiarthroplasty, conversion to, for treatment of hip instability, 605
Herbert technique, for distal radial ulnar joint replacement, 369–370
Hip abductors, repair of, for treatment of hip instability, 607
Hip dislocation. See *Dislocation of hip*.
Hip instability, treatment of, **593–610**
assessment of cause of instability, 596
closed reduction as definitive treatment, 597–598
examination, 595
final outcome, 608
history, 594–595
initial management, 596–597
open reduction, 598
radiologic assessment, 595–596
revision arthroplasty, 598
subluxation vs. dislocation, 593–594
surgical treatment options, 598–599
acetabular component repositioning, 599–600
change of acetabular implants design, 600–603
conversion to hemiarthroplasty, 605
femoral component reorientation, 603
increasing abductor tension, 606
increasing head size, 604–605
increasing neck length, 606
increasing offset of height of femoral component, 606
reattachment of displaced nonunion of trochanteric osteotomy, 606
removing impingement, 605–606
soft tissue procedures, 607–608
trochanteric advancement, 606–607

Iliac resection for tumor, instability and, 698
Impingement, removing, for treatment of hip instability, 605–606
Instability after major joint replacement, axial instability of knee, **627–637**
dislocation after total hip arthroplasty, **587–591**
epidemiology of instability after major hip replacement, **543–552**
experimental and computational simulation of total hip arthroplasty dislocation, **553–567**
instability after major tumor resection, **697–710**
instability after prosthetic arthroplasty of shoulder, **649–659**
instability after total elbow arthroplasty, **671–677**
instability of shoulder arthroplasty, **661–670**
issue on, **543–722**
preoperative planning to prevent dislocation of hip, **579–586**
preoperative planning to prevent instability in total knee arthroplasty, **611–626**
rotational instability and malrotation after total knee arthroplasty, **639–647**
treatment of hip instability, **593–610**
treatment of instability after total elbow arthroplasty, **679–695**
virtual reality computer animation of effect of component position and design on stability after total hip arthroplasty, **569–577**
Internal impingement, in professional baseball players, 443
Interphalangeal dislocations, 8–9
Ischiopubic resection for tumor, instability and, 702–703

Jones fractures, 4, 6. See also *Metatarsal(s), fifth, base of, fractures of*.

Kienböck's disease, vascularized bone grafts for, **263–278**
historical perspective, 268
indications and contraindications, 270
technique of harvest, 272–275
Kirschner wire fixation, of tarsometatarsal joint injuries, 17
Knee, axial instability of, **627–637**
flexion instability, 632–634
varus-valgus instability, 627–629
results of correction, 631–632
soft tissue balancing in total knee arthroplasty, 629–631
Knot, "new" arthroscopic sliding, **523–524**

Ligament laxity and prosthetic constraint, for preventing instability in total knee arthroplasty, 619
Ligament reconstruction, for treatment of instability after total elbow arthroplasty, 689, 691
Lisfranc fractures, in multiply injured patient, 199
Lisfranc's level amputations, muscle balancing after, 186

Maisonneuve fractures, 85–86
Major hip replacement. See also *Total hip arthroplasty*.
epidemiology of instability after, **543–552**
prevalence, 543–544

risk factors for initial episode of dislocation, factors related to implant design, 548–550
factors related to patient, 544–545
factors related to surgical technique, 545–548
risk factors for late dislocation, 550
risk factors for recurrent dislocation, 550
Malleolus fractures, isolated lateral, management of, 84
isolated medial, management of, 84–85
lateral, fixation of, 87–88
medial, fixation of, 88
posterior, fixation of, 88
Metatarsal(s), base of, fracture of, 2
fifth, anatomy, 173
base of, fractures of, **171–180**
acute Jones fracture (type II), 175
history, 171–173
postoperative management, 177
summary, 178–179
surgical technique, 175–177
treatment of diaphyseal stress fractures (type III) and delayed unions and nonunions, 177–178
tuberosity avulsion fractures (type I), 174–175
blood supply, 173–174
fracture of, 4
fractures of, 1–2
in multiply injured patient, 199
injuries to toes and, **1–10**
neck, fracture of, 3–4
shaft, fracture of, 2–3
Metatarsophalangeal joint, first, dislocation of, 7–8
injuries to, dislocation of first metatarsophalangeal joint, 7–8
fracture of phalanges, 8
interphalangeal dislocations, 8–9
turf toe, 6–7
Microsurgical free flaps, for soft tissue reconstruction for calcaneal fractures or osteomyelitis, 158, 165
Midfoot, anatomy of, 21–22
combined injuries of, 30–32
fractures of, in multiply injured patient, 199
use of ring fixators in, 212
Monopolar electrothermal arthroscopy for treatment of shoulder instability in athlete, **511–521**
applications of thermal stabilization of shoulder, 514
clinical results, 518–519
complications and failures, 519–520
methods of thermal energy delivery, 512–514
postoperative care, multidirectional instability, 518
posterior instability, 517–518
unidirectional anterior instability, 517
scientific background, 511–512
surgical technique, 514–517
Multidirectional shoulder instability, treatment of athlete with, **475–484**
anterior approach, 477–480
epidemiology, 475
evaluation, 476–477
nonoperative, 477
operative, 477
pathophysiology, 476
rehabilitation, 481–482
results, 482–483
thermal capsular shifting procedure, 480–481
Multiply injured patients, outcome of foot injuries in, **193–204**
authors' review, 199–202
compartment syndrome, 196
general treatment principles, 195–197
high-energy injuries, 194–195
initial management of trauma patient, 193–194
investigations, 197–198
missed or delayed diagnosis, 194
reasons for, 202
specific injuries, calcaneus, 198–199
Lisfranc, 199
metatarsal, 199
midfoot, 199
talus, 198
timing of surgery, 197
vascular compromise, 196
Muscle balancing, after foot and ankle amputations, 186

Navicular bone, avulsion fractures of, 25
body fractures of, 25–27
stress fractures of, 28–29
Neck length, increasing, for treatment of hip instability, 606
Nerve injuries about shoulder, in athletes, 395–396
Nerve transection, foot and ankle, salvage vs. amputation for, 182
Neuroarthropathic (Charcot) fractures, use of ring fixators in, 209

Open reduction, for treatment of hip instability, 598
Osteoarticular allografts, for proximal femur reconstruction, 704–705
Osteochondral allograft, for distal femur reconstruction, 705–706
for proximal tibia reconstruction, 707–708
Osteoinductive materials, as bone graft substitute for unstable distal radius fractures, 349
Osteomyelitis, calcaneal fractures or, soft tissue reconstruction for. See *Calcaneal fractures, or osteomyelitis, soft tissue reconstruction for.*
Osteopenic bone, in diabetic patients, fixation for, 129
Overhead athlete, biceps lesions in, 488–492
coracoid impingement in, 485–486
coracoid impingement syndrome, rotator interval reconstruction, and biceps tenodesis in, **485–493**
evaluation and management of shoulder instability in, **423–430**
biomechanics of throwing, 424
epidemiology, 424
history, 425
physical examination, 425–426
postoperative management, 429–430
radiology, 426
rehabilitation, 426–429
shoulder pathophysiology, 424–425
rotator interval lesions, 486, 488
SLAP lesions in, **431–441**
acceleration vs. deceleration injury and proposed mechanism, 436–437
arthroscopic evaluation and treatment, 437–438
arthroscopic repair and rehabilitation, 438
biomechanics, 434–436
conclusions, 438–440
evaluation, 437
historical perspective, 431–432
peel-back sign, 432–433
shoulder at risk: 180° rule, 438

Patellectomy, instability in total knee arthroplasty and, 624
Patellofemoral joint, instability after major tumor resection in, 708
instability in total knee arthroplasty and, 624–625
Peel-back sign, in overhead athlete with SLAP lesions, 432–433
Pelvic resection for tumor, instability and, 697–703
Percutaneous fixation, dorsal, technique of, for scaphoid fractures, arthroscopy, 251
cannulated screw placement, 251–252
guidewire placement, 250–251
overview, 248
postoperative care, 252–253
setup, 248–249
history of, for scaphoid fractures, 255
Periacetabular resections, instability and, 698–702
Phalanges, fracture of, 8
Pilon fractures, authors' preferred technique for management of, 98–100
classification, 91–93
external fixation alone as definitive fixation, 100
hybrid external fixation accompanied by percutaneous fixation, 100–101
radiographic evaluation, 93–94
summary, 101
treatment of, **91–102**
nonoperative, 94
operative, 94–97
timing of, 97–98
Posterior cruciate ligament retention, for preventing instability in total knee arthroplasty, 620
Posterior instability of glenohumeral joint, **463–473**
authors' preferred operative technique, 467–470
discussion, 471–472
evaluation, 464–465
pathophysiology, 463–464
rehabilitation, 470
results, 470–471
treatment, 465–467
Prosthetic arthroplasty of shoulder, instability after, **649–659**
classification, 649
evaluation, 652–656
review of literature, anterior instability, 650
inferior instability, 651
posterior instability, 650–651
superior instability, 650
treatment, 656–658
Proximal femur, instability after major tumor resection in, 703–705
Proximal tibia, instability after major tumor resection in, 707–708

Radial head component, addition of, for treatment of instability after total elbow arthroplasty, 691–692
Recurvatum, correcting, for preventing instability in total knee arthroplasty, 623
Rehabilitation, after multidirectional shoulder instability, 481–482
shoulder, strategies, guidelines, and practice, basic science, 525–527
guidelines, complete and accurate diagnoses, 527
glenohumeral rehabilitation, 531–532
for progression, 533–534
proximal segment control, 527, 529
pylometric exercises, 532–533
for return to play, 535
scapular rehabilitation, 529–530
practice, 527, 529–533
for shoulder injuries in elite thrower, 426–429
after surgery for posterior instability of glenohumeral joint, 470
Resection arthroplasty, for failed total elbow arthroplasty, 694
Revision arthroplasty, for treatment of hip instability, 598
Ring fixators, applications of, in complex foot and ankle trauma, **205–214**
conclusions, 212
historical perspective, 205
reasons for using, 205–206
specific fractures, calcaneal fractures, 210–212
foot fractures, 212
neuroarthropathic (Charcot) fractures, 209
talus fractures, 209–210
tibial plafond fractures, 206–209
Rotational instability, after total knee arthroplasty, 639–640
Rotator cuff repairs, all arthroscopic, **501–510**
discussion, 509–510
full-thickness tears, acromioclavicular joint, 504
acromioplasty, 504
anchor placement, 505
anchor selection, 505
cuff mobilization, 503–504
instruments, 508
knot tying, 507
repair site preparation, 504–505
suture placement and passing, 505, 507
suture selection, 505
tear classification, 503
visualization, 503
indications, 501
operative technique, inspection of glenohumeral joint, 502–503
portals, 502
positioning, 501–502
partial-thickness tears, 503
postoperative management, 508
results, 508–509
Rotator cuff tears, anterior superior instability with, **457–461**
clinical examination, 457–458
diagnostic studies, 458
discussion, 459–461
materials and methods, 457
results, 459
surgical findings, 458–459
in athletes, 395
partial-thickness, in professional baseball players. See *Arthroscopic repair of partial-thickness rotator cuff tears and SLAP lesions in professional baseball players.*
Rotator cuff tendonitis, in athletes, 397
Rotator interval lesions, in overhead athlete, aftercare, 488
anatomy and pathophysiology, 486
clinical presentation, physical examination, and diagnosis, 486, 488
surgical technique, 488
treatment, 488

Saddle prosthesis, for pelvic reconstruction, 699–700

Salvage, after complex foot and ankle trauma. See *Foot and ankle trauma, complex, salvage or amputation after.*
Scaphoid, anatomy of, 233–234
fractures, arthroscopic reduction and percutaneous fixation of, with a novel dorsal technique, **247–261**
history of percutaneous and arthroscopic fixation of, clinical results, 256–259
open techniques, 255
percutaneous fixation, 255
use of arthroscopy, 255–256
indications for fixation of, 253
principles of fixation of, mechanics, 253–254
soft tissue management, 254
vascularity, 254–255
summary, 259
technique of dorsal percutaneous fixation of, arthroscopy, 251
cannulated screw placement, 251–252
guidewire placement, 250–251
overview, 248
postoperative care, 252–253
setup, 248–249
instability, dynamic, arthroscopic and open management of, **233–240**
arthroscopic treatment, 236–239
diagnosis, 234
diagnostic arthroscopy, 234–235
management, 234
open management, 239
summary, 239
nonunion, vascularized bone grafts for, **263–278**
historical perspective, 267–268
indications and contraindications, 269–270
technique of harvest, 270–272
Scapholunate ligament injuries, bone-retinaculum-bone reconstruction of, **241–246**
discussion, 244–245
indications, 241–242
outcomes, 244
postoperative care, 243–244
surgical technique, 242–243
Scapula fractures, in athletes, 394
Scheker technique, for distal radial ulnar joint replacement, 370–373
Screw fixation, of tarsometatarsal joint injuries, 17
Shoulder arthroplasty, instability of, **661–670**
factors determining mobility after glenohumeral arthroplasty, 662–663
factors determining stability after glenohumeral arthroplasty, 663–664
unstable glenohumeral arthroplasty, clinical syndromes, 664–666
evaluation, 666
management and prevention, 666–668
prosthetic, instability after, **649–659**
classification, 649
evaluation, 652–656
review of literature, anterior instability, 650
inferior instability, 651
posterior instability, 650–651
superior instability, 650
treatment, 656–658
Shoulder injuries, in sports, acute, 393–396
chronic, 396–397
differential diagnosis of, **393–398**
Shoulder instability, anatomy, pathophysiology, and biomechanics of, **381–391**
anterior superior, with rotator cuff tearing: SLAC lesion, **457–461**
clinical examination, 457–458
diagnostic studies, 458
discussion, 459–461
materials and methods, 457
results, 459
surgical findings, 458–459
atraumatic, 388
basic science, 381–383
clinical examination, 388–389
evaluation and management of, in elite overhead thrower, **423–430**
biomechanics of throwing, 424
epidemiology, 424
history, 425
physical examination, 425–426
postoperative management, 429–430
radiology, 426
rehabilitation, 426–429
shoulder pathophysiology, 424–425
microtraumatic, 386–388
monopolar electrothermal arthroscopy for treatment of, in athlete, **511–521**
applications of thermal stabilization of shoulder, 514
clinical results, 518–519
complications and failures, 519–520
methods of thermal energy delivery, 512–514
postoperative care, multidirectional instability, 518
posterior instability, 517–518
unidirectional anterior instability, 517
scientific background, 511–512
surgical technique, 514–517
multidirectional, treatment of athlete with, **475–484**
anterior approach, 477–480
epidemiology, 475
evaluation, 476–477
nonoperative, 477
operative, 477
pathophysiology, 476
rehabilitation, 481–482
results, 482–483
thermal capsular shifting procedure, 480–481
posterior, **463–473**
authors' preferred operative technique, 467–470
discussion, 471–472
evaluation, 464–465
pathophysiology, 463–464
rehabilitation, 470
results, 470–471
treatment, 465–467
in professional baseball players, 444
traumatic, 383–384
anterior translation, 384–386
inferior translation, 386
posterior translation, 386
traumatic anterior, **399–409**
in athletes, arthroscopic Bankart repair with Suretac device for, **411–421**
anatomy of shoulder stability, 411–412
complications, 418
discussion, 418–419
patient selection, 412–413
postoperative care, 415–416
results, 416–417
surgical technique, 413–415
thermal capsulorrhaphy, 416

Shoulder instability (*Continued*)
incidence, 399–400
persistent, 408
results, 408
surgical stabilization, 400–401
suture anchor technique for arthroscopic stabilization, 401
diagnostic arthroscopic evaluation, 401–403
glenoid preparation, 403–404
postoperative care, 406–408
suture anchor placement and Bankart repair, 404–406
Shoulder pain, anterior, causes of, 396
lateral, causes of, 396–397
posterior, causes of, 397
superior, causes of, 397
Shoulder rehabilitation strategies, guidelines, and practice, **525–536**
basic science, 525–527
guidelines, complete and accurate diagnoses, 527
glenohumeral rehabilitation, 531–532
for progression, 533–534
proximal segment control, 527, 529
pylometric exercises, 532–533
for return to play, 535
scapular rehabilitation, 529–530
practice, 527, 529–533
Simulation, experimental and computational, of total hip arthroplasty dislocation, **553–567**
discussion, 563–566
materials and methods, computational simulation, 557–559
experimental simulation, 554–557
results, 559–563
Skin graft, for acute or chronic lateral calcaneal wound, 153
SLAC lesion, **457–461**
clinical examination, 457–458
diagnostic studies, 458
discussion, 459–461
materials and methods, 457
results, 459
surgical findings, 458–459
SLAP lesions, in overhead athlete, **431–441**
acceleration vs. deceleration injury and proposed mechanism, 436–437
arthroscopic evaluation and treatment, 437–438
arthroscopic repair and rehabilitation, 438
biomechanics, 434–436
conclusions, 438–440
evaluation, 437
historical perspective, 431–432
peel-back sign, 432–433
shoulder at risk: 180° rule, 438
in professional baseball players. See *Arthroscopic repair of partial-thickness rotator cuff tears and SLAP lesions in professional baseball players.*
Soft tissue injuries, in athletes, 394
foot and ankle, salvage vs. amputation for, 181–182
Soft tissue reconstruction, for calcaneal fractures or osteomyelitis. See *Calcaneal fractures, or osteomyelitis, soft tissue reconstruction for.*
Soft tissue repair, for treatment of instability after total elbow arthroplasty, 689
Subacromial impingement, in professional baseball players, 445
Subluxation vs. dislocation, in hip instability, 593–594
Subscapularis tendon, anatomy, 495
authors' preferred postoperative treatment protocol, 499
clinical presentation, 496
imaging, 496–497
mechanism of injury, 496
results, 499
technique for repair of, **495–500**
authors' preferred method, 497–499
Superior labrum repair, in professional baseball players, 448–450
Suretac device. See *Arthroscopic Bankart repair with Suretac device for traumatic anterior shoulder instability in athletes.*
Suture anchor technique, for arthroscopic stabilization of anterior shoulder instability, 401–408
Swanson technique, for distal radial ulnar joint replacement, 367–369
Syndesmosis injuries, 85–86

Talar body fractures, **65–77**
comminuted body fractures, 75–76
fractures of lateral process, 74–75
clinical presentation and treatment, 75
fractures of posterior tuberosity, clinical presentation and treatment, 74
osteochondral fractures of talar dome, clinical presentation, 65
radiographic evaluation and classification, 65–67
treatment of, 67–68
operative, 68
preferred method of, 68–69
shear fractures of talar body, author's preferred operative technique, 71–72
clinical presentation, 70
complications, 72, 74
radiographic evaluation, 70
treatment, 71
Talar neck fractures, **53–64**
classification, 57–58
clinical presentation, 56
complications, 61–62
history, 53
mechanism of injury, 55–56
in multiply injured patient, 198
operative management, 59–61
postoperative care and rehabilitation, 61
radiographic evaluation, 56–57
summary, 62–63
treatment, group I, 58
group II, 58–59
group III, 59
group IV, 59
Talus, anatomy, 53–55
Talus fractures, use of ring fixators in, 209–210
Tarsal bones, fractures of, **21–33**
conclusion, 32
cuboid bone, 22–25
cuneiform bones, 29–32
midfoot anatomy, 21–22
navicular bone, 25–29
Tarsometatarsal joint injuries, classification, 11–12
conclusion, 19
mechanism, 12–13
principles of treatment, nonoperative, 16
operative, 16–19

radiographic evaluation, 13–16
recent developments and advances in diagnosis and treatment, **11–20**
Thermal capsular shifting procedure, for multidirectional shoulder instability, 480–481
Thermal capsulorrhaphy, for reducing capsular laxity, 416
Throwing, biomechanics of, 424
Throwing athlete. See *Overhead athlete.*
Tibia, proximal, instability after major tumor resection in, 707–708
resection of, for preventing instability in total knee arthroplasty, 614–616
Tibial plafond fractures, use of ring fixators in, 206–209
Toe(s), amputations, muscle balancing after, 186
and metatarsals, injuries to, **1–10**
Total elbow arthroplasty, instability after, **671–677**
anatomy, 673–674, 680–681
cause, implant design, 682, 685
patient, 688
postoperative management, 688
surgeon, 688
surgical approach, 685–688
classification, 681–682
evaluation, 688–689
future considerations, 694
implications for linked total elbow arthroplasties, 694
magnitude and relevance of problem, 680
operative technique, 675
patient selection, 674–675
postoperative care, 675–676
problem, 671–673
relevant biomechanics, 680–681
treatment of, **679–695**
addition of radial head component, 691–692
closed reduction and cast, 689
component revision, 692–693
ligament reconstruction, 689, 691
revision to linked total elbow arthroplasty, 693–694
soft tissue repair, 689
unstable nonconstrained, treatment of, 676
Total femoral replacement for tumor, instability after, 708–709
Total hip arthroplasty. See also *Major hip replacement.*
dislocation after, **587–591**
discussion, 590–591
materials and methods, 587
preoperative planning to prevent, 579–586
decreasing incidence of dislocation, component design and size, 582–582
component positioning, 583–584
operative factors, 581
patient risk factors, 579–581
postoperative treatment, 584
results, 587–590
stability after, virtual reality computer animation of effect of component position and design on, **569–577**
materials and methods, 569–572
results, 572–576
Total hip arthroplasty dislocation, experimental and computational simulation of, **553–567**
discussion, 563–566
materials and methods, computational simulation, 557–559
experimental simulation, 554–557
results, 559–563
Total knee arthroplasty, preoperative planning to prevent instability in, **611–626**
anterior or posterior instability, 619–620
balancing flexion and extension gap, 620–621
clinical examination, 611–612
general principles for achieving balanced knee arthroplasty and how to prevent instability, femoral bone resection, 616–619
tibial resection, 614–616
history, 611
joint line level, 621
ligament laxity and prosthetic constraint, 619
overall limb alignment, 621–623
patellofemoral instability, 624–625
posterior cruciate ligament retention, 620
radiographic examination, 612
special cases, 623–624
surgical approach, 612–614
rotational instability and malrotation after, **639–647**
analyzing result of computed tomography rotation, 644
discussion, 644–646
intraoperatively determining femoral and tibial component rotation during primary or revision total knee arthroplasty, 640–641
preoperatively determining femoral and tibial component rotation in malfunctioning total knee arthroplasty, 641–644
rotational instability, 639–640
Transmetatarsal amputations, muscle balancing after, 186
Trauma, foot and ankle. See *Foot and ankle trauma.*
wrist, advanced techniques in management of, issue on, **219–379**
Trauma patient. See also *Multiply injured patients.*
initial management, 193–194
Triangular fibrocartilage complex injuries, anatomy, 279–281
arthroscopic repair of, **279–294**
for irreparable lesions, 291–292
postoperative care, 292
for reparable radial-sided lesions, 289–291
for reparable ulnar-sided lesions, 286, 288–289
biomechanics, 281
classification, 284–286
diagnosis, diagnostic modalities, 283–284
differential diagnosis, 282
history, 282
physical examination, 282
results of repair of, 292
summary, 292–293
treatment, nonoperative, 286
open surgical repair, 286, 287
ulnar recession, 286
Trimalleolar fractures, 85
Trochanteric advancement, for treatment of hip instability, 606–607
Trochanteric osteotomy, reattachment of displaced nonunion of, for treatment of hip instability, 606
Tumor resection, major, instability after, **697–710**
distal femur, 705–707
patellofemoral joint, 708
pelvic resection, 697–698
iliac resection, 698
ischiopubic resection, 702–703
periacetabular resections, 698–702
proximal femur, 703–705
proximal tibia, 707–708
total femoral replacement, 708–709
Turf toe, 6–7

uHead prosthesis, for distal radial ulnar joint replacement, 370

Varus-valgus instability, of knee, 627–629
results of correction, 631–632
soft tissue balancing in total knee arthroplasty, 629–631
Varus-valgus stability, for preventing instability in total knee arthroplasty, 621–622
Vascular compromise, from foot injuries in multiply injured patient, 196
Vascular injury, foot and ankle, salvage vs. amputation for, 182
Vascularized bone grafts, for scaphoid nonunion and Kienböck's disease, **263–278**
experimental studies, 268–269
historical perspective, 267–268
pedicled vascularized bone grafts from dorsal distal radius, indications and contraindications, 269–270
technique of harvest, 270–275
principles, 263
summary, 275, 277
vascular anatomy of dorsal distal radius and ulna, 264–267
Virtual reality computer animation of effect of component position and design on stability after total hip arthroplasty, **569–577**
materials and methods, 569–572
results, 572–576

Wound healing, in diabetic patients with ankle fractures, 116–119
Wrist trauma, advanced techniques in management of, issue on, **219–379**